



Literaturverzeichnis zu

Dr. Andrea Benecke Psychotherapie bei Diabetes

Psychotherapie Aktuell
Heft 2.2009
ISSN 1869-0335

Affenito SG, Bachstrand JR, Welch GW, et al. (1997). Subclinical and Clinical Eating Disorders in IDDM Negatively Affect Metabolic Control. *Diabetes Care*, 20, 2, 182-184.

Anderson RJ, Freedland KE, Clouse RE, Lustman PJ (2001). The Prevalence of Comorbid Depression in Adults With Diabetes. A meta-analysis. *Diabetes Care*, 24, 6, 1069-1078.

Arnold-Wörner N, Holle R, Rathmann W, Mielck A (2008). The importance of specialist treatment and diabetes education für the compliance of subjects with type 2 diabetes – results from a population based survey. *Exp Clin Endocrinol Diabetes*, 116 (2), 123-28.

Black SA, Markides KS, Bray LA (2003). Depression predicts increased incidents of adverse health outcomes in older Mexican Americans with type 2 diabetes. *Diabetes Care*, 26 (10):2822-2828.

Cochran J & Conn VS (2008). Meta-analysis of Quality of Life Outcomes Following Diabetes Self-management Training. *The Diabetes Educator*; 34:815-823.

Cox DJ, Gonder-Frederick L, Polonsky W et al. (2001). Blood Glucose Awareness Training (BGAT-2). *Diabetes Care*, 24, 4:637-642.

Derijks HJ, Meyboom RH, Heerdink ER, et al. (2008). The association between antidepressant use and disturbances in glucose homeostasis: Evidence from spontaneous reports. *Eur J Clin Pharmacol*, 64 (5):531-538.

Diabetes Control and Complications Trial Research Group (1993). The Effect of Intensive Treatment of Diabetes on the Development and Progression of Long-Term Complications in Insulin-Dependent Diabetes Mellitus. *New England Journal of Medicine*, 329, 14:977-986.

Glasgow R, Ruggiero L, Eakin E, et al. (1997). Quality of Life and Associated Characteristics in a Large National Sample of Adults With Diabetes. *Diabetes Care*, 20,4:562-567.

Goebel-Fabbri AE, Fikkan J, Franko DL et al (2008). Insulin restriction and associated morbidity and mortality in women with Type 1 diabetes. *Diabetes Care*, 31 (3):415-419.

Grey M, Boland EA, Davidson M et al. (2000). Coping skills training for youths with diabetes mellitus has long-lasting effects on metabolic control and quality of life. *The Journal of Pediatrics*, 137 (1):107-113.



- Hayes AJ, Clarke PM, Glasziou PG et al. (2008). Can self-rated health scores be used for risk prediction in patients with type 2 diabetes? *Diabetes Care*, 31 (4):795-797.
- Ismail K, Winkley K, Rabe-Hesketh S (2004). Systematic review and meta-analysis of randomised controlled trials of psychological interventions to improve glycaemic control in patients with type 2 diabetes. *The Lancet*, 363, May 15:1589-1597.
- Kleefstra N, Landman GW, Houweling ST et al. (2008). Prediction of mortality in type 2 diabetes from health-related quality of life (ZODIAC-4). *Diabetes Care*, 31(5):932-33.
- Li C, Barker L, Ford ES et al. (2008). Diabetes and anxiety in US adults: Findings from the 2006 Behavioral Risk Factor Surveillance System. *Diabetic Medicine* 25 (7):878-881.
- Maddigan SL, Feeny DH, Johnson JA (2005). Health-related quality of life deficits associated with diabetes and comorbidities in a Canadian National Population Health Survey. *Quality of Life Research* 14:1311-1320.
- Olmsted MP, Colton PA, Daneman D et al. (2008). Prediction of the onset of disturbed eating behavior in adolescent girls with type 1 diabetes. *Diabetes Care*, 31 (10):1978-1982.
- Saito I, Inami F, Ikebe T, et al. (2006). Impact of diabetes on healthrelated quality of life in a population study in Japan. *Diabetes Research and Clinical Practice*, 73:51-57.
- Snoek FJ, van der Ven NCW, Lubach CHC et al.(2001). Effects of cognitive behavioural group training (CBGT) in adult patients with poorly controlled insulin-dependent (type 1) diabetes: a pilot study. *Patient Education and Counseling*, 45:143-148.
- Varni JW, Limbers CA, Burwinkle TM et al. (2008). The ePedsQL in type 1 and type 2 diabetes: feasibility, reliability, and validity of the Pediatric Quality of Life Inventory Internet administration. *Diabetes Care* 31 (4):672-677.
- Winkley K, Landau S, Eisler I, Ismail K (2006). Psychological interventions to improve glycaemic control in patients with type 1 diabetes: systematic review and meta-analysis of randomised controlled trials. *British Medical Journal*, 333 (7558):65.

Dr. Andrea Benecke
Psychologisches Institut der Universität Mainz, Abteilung klinische Psychologie
Wallstr. 3
55122 Mainz
benecke@uni-mainz.de





Literaturverzeichnis zu

Dr. Hans-Günter Budde **Psychotherapie bei Herz-Kreislauf- Erkrankungen**

Psychotherapie Aktuell
Heft 2.2009
ISSN 1869-0335

- Appels, A., Mulder, P. (1989) Fatigue and heart disease: The association between „vital exhaustion“ and past, present and future coronary heart disease. *Journal of Psychosomatic Research*, 33, 727-738.
- Bardé, B., Jordan, J. (2003) *Psychodynamische Beiträge zu Ätiologie, Verlauf und Psychotherapie der koronaren Herzkrankheit*. Frankfurt / Main: VAS.
- Barth, J., Bengel, J. (2003) *Interventionen zur Raucherentwöhnung bei kardiovaskulären Erkrankungen*. Frankfurt / Main: VAS.
- Bart, J., Schumacher, M., Herrmann-Lingen, Ch. (2004) Depression as a risk factor for mortality in patients with coronary heart disease: a meta-analysis. *Psychosom. Medicine* 66, 645-650.
- Blumenthal, J.A. et al. (1997) Stress management and exercise training in cardiac patients with myocardial ischemia. *Archives of Internal Medicine*, 157, 2213-2223.
- Blumenthal, J.A. et al. (2002) Usefulness of psychosocial treatment of mental stress induced myocardial ischemia in men. *American Journal of Cardiology* 89, 164-168.
- Budde, H.G. (2006) Psychologische Faktoren oder Verhaltensfaktoren bei andernorts klassifizierten Erkrankungen – Cardiologie: Herz-Kreislauf-Erkrankungen. In Beiglböck, W., Feselmayer, S. & Honemann, E. (Hrsg.). *Handbuch der klinisch-psychologischen Behandlung*. Wien / New York: Springer, 443 - 464.
- Budde, H.G., Langosch, W. (2004) Psychologische Interventionen bei koronarer Herzkrankheit (KHK) – Empirische Befunde und Schlussfolgerungen für die praktische Versorgung. *Psychotherapeutische Praxis*, 4, 177 - 182.
- Cossette, S., Frasure-Smith, N., Lespérance, F. (2001) Clinical implications of a reduction in psychological distress on cardiac prognosis in patients participating in a psychological intervention program. *Psychosomatic Medicine*, 63, 257-266.
- Denollet, J. (2000) Type D personality: A potential risk factor redefined. *Journal of Psychosomatic Research*, 49, 255-266.
- Dusseldorp, E. et al. (1999) A meta-analysis of psychoeducational programs for coronary heart disease patients. *Health Psychology*, 18, 506-519.
- Frasure-Smith, N. et al. (1997) Randomised trial of home-based psychosocial nursing intervention for patients recovering from myocardial infarction. *Lancet*, 350, 473-479.

- Frasure-Smith, N., Prince, R.H. (1985) The Ischemic Heart Disease Life Stress Monitoring Program: Impact on mortality. *Psychosomatic Medicine*, 47, 431-445.
- Frasure-Smith, N., Prince, R.H. (1989) Long-term follow up of the Ischemic Heart Disease Life Stress Monitoring Program. *Psychosomatic Medicine*, 51, 431-445.
- Grande, G. (2008) Genderspezifische Aspekte der Gesundheitsversorgung und Rehabilitation nach Herzinfarkt. *Bundesgesundheitsblatt – Gesundheitsforschung – Gesundheitsschutz*, 51, 36-45.
- Grande, G., Badura, B. (2001) Die Rehabilitation der KHK aus gesundheitssystemanalytischer Perspektive. Frankfurt / Main: VAS.
- Hank, P., Mittag, O. (2003) Zur Bedeutung von Ärger und Ärgerausdruck für die Entstehung und Prognose der koronaren Herzkrankheit. Frankfurt / Main: VAS.
- Herrmann-Lingen, C., Buss, U. (2002) Angst und Depressivität im Verlauf der koronaren Herzkrankheit. Frankfurt / Main: VAS.
- Herrmann-Lingen, C., Albus, C., Titscher, G. (Hrsg.) (2008) *Psychokardiologie*. Köln: Deutscher Ärzte-Verlag.
- Jones, D.A., West, R.R. (1996) Psychological rehabilitation after myocardial infarction: Multicentre randomised control trial. *British Medical Journal*, 313, 1517-1521.
- Ladwig, K.-H., Erazo, N., Rugulies, R. (2004) Zur Bedeutung von Ärger und Ärgerausdruck für die Entstehung und Prognose der koronaren Herzkrankheit. Frankfurt / Main: VAS.
- Laederach-Hofmann, K., Messerli-Buergy, N. (2004) Panikstörung, kardiales Syndrom X und koronare Herzkrankheit. Frankfurt / Main: VAS.
- Langosch, W., Budde, H.G., Hahn, P. (2004) Psychosomatik des Herzinfarktes. In Roskamm, H., Neumann, F.-J., Kalusche, D. & Besthorn, H.-P. (Hrsg.) *Herzkrankheiten – Pathophysiologie, Diagnostik, Therapie*. 5. Aufl.; Berlin, Heidelberg, New York: Springer, 1269 – 1283.
- Langosch, W., Budde, H.G., Linden, W. (2003) Psychologische Interventionen zur koronaren Herzkrankheit (Stress-Bewältigung, Entspannungsverfahren, Ornish-Gruppen). Frankfurt / Main: VAS.
- Langosch, W., Budde, H.G., Linden, W. (2007) Psychological Interventions for Coronary Heart Disease: Stress Management, Relaxation, and Ornish Groups. In Jordan, J., Barde, B. & Zeiher, A. (Hrsg.) *Contributions Toward Evidence-Based Psychocardiology*. New York: APA, 231 – 254.
- Lazanowski, C., Jordan, J. (2003) Psychosoziale Aspekte der Herzkatheteruntersuchung, Koronarangiographie und –angioplastie (PTCA). Frankfurt / Main: VAS.
- Linden, W. (2000) Psychological treatments in cardiac rehabilitation: Review of rationals and outcomes. *Journal of Psychosomatic Research*, 48, 443-454.
- Linden, W., Stossel, C., Maurice, J. (1996) Psychosocial interventions for patients with coronary artery disease. *Archives of Internal Medicine*, 256, 745-752.
- Löwel, H., Heier, M., Hörmann, A., Meisinger, C. (2005) Koronare Herzkrankheiten bei Männern und Frauen. *Praxis Verhaltenstherapie. Klin. Verhaltensmed. Rehab.* 18, 226-231.



- Myrtek, M. (2000) Das Typ-A-Verhaltensmuster und Hostilität als eigenständige Risikofaktoren der koronaren Herzkrankheit. Frankfurt / Main: VAS.
- Myrtek, M. (2001) Meta-analyses of prospective studies on coronary heart disease, Type A personality, and hostility. *International Journal of Cardiology*, 79, 245-251.
- Ong, L., Linden, W., Young, S. (2004) Stress management: What is it? *Journal of Psychosom. Research* 56, 133-137.
- Ornish, D. (1991) *Open Your Heart*. New York: Random House. / (1992) *Revolution in der Herztherapie*. Stuttgart: Kreuz.
- Ornish, D. (1998) Avoiding revascularization with lifestyle changes: The Multicenter Lifestyle Demonstration Project. *American Journal of Cardiology* 82, 72T-76T.
- Ornish, D. (1999) *Die revolutionäre Therapie: Heilen mit Liebe*. München: Mosaik-Verlag.
- Ornish, D. et al. (1990) Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet* 336, 129-133.
- Ornish, D. et al. (1998) Intensive lifestyle changes for reversal of coronary heart disease. *JAMA*, 280, 2001-2007.
- Orth-Gomér, K. et al. (2009) Stress Reduction Prolongs Life in Women with Coronary Disease – The Stockholm Women’s Intervention Trial For Coronary Heart Disease (SWITCHD). *Circulation: Cardiovascular Quality and Outcomes*, 2, 25-32.
- Rees, K., Bennett, P., West, R., Davey Smith, G., Ebrahim, S. (2004) Psychological interventions for coronary heart disease. *Cochrane Database of Systematic Reviews*, Issue 2. Art.No.: CD002902. DOI: 10.1002/14651858.CD002902.pub2.
- Rosengreen, A. et al. (2004) Association of psychosocial risk factors with risk of acute myocardial infarction in 11119 cases and 13648 controls from 52 countries (the INTERHEART study). *Lancet* 364, 953-962.
- Rozanski, A., Blumenthal, J.A., Kaplan, J. (1999) Impact of psychological factors on the pathogenesis of cardiovascular disease and implications for therapy. *Circulation*, 99, 2192-2217.
- Rugulies, R. (2002) Depression as a predictor for coronary heart disease: A review and meta-analysis. *American Journal of Preventive Medicine*, 23, 51-61.
- Rugulies, R., Siegrist, J. (2007) Sociological Aspects of the Development and Course of Coronary Heart Disease: Social Inequality and Chronic Emotional Distress in The Workplace. In Jordan, J., Barde, B. & Zeiger, A. (Hrsg.) *Contributions Toward Evidence-Based Psychocardiology*. New York: APA, 13 – 33.
- Schlicht, W., Kanning, M., Bös, K. (2003) *Psychosoziale Interventionen zur Beeinflussung des Risikofaktors Bewegungsmangel*. Frankfurt / Main: VAS.
- Sheps, D.S., Soufer, R., Freedland, K.E. (2007) Psychological stress and myocardial ischemia : Understanding the link and the implications. *Psychosom. Medic.* 69, 491-492.
- Skala, J.A., Freedland, K.E., Carney, R.M. (2005) *Heart Disease*. Göttingen: Hogrefe & Huber.





Titscher, G., Schöppl, Ch. (2000) Die Bedeutung der Paarbeziehung für Genese und Verlauf der koronaren Herzkrankheit. Frankfurt / Main: VAS.

Weidemann, H., Meyer, K., Fischer, T., Wetzell, A. (2003) Frauen und koronare Herzkrankheit. Frankfurt / Main: VAS.

Writing Committee for the ENRICHD investigators (2003) Effects of treating depression and low perceived social support on clinical events after myocardial infarction: The Enhancing Recovery in Coronary Heart Disease patients (ENRICHD) randomized trial. JAMA 289, 3106-3116.

Yusuf, S. et al. (2004) Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study). Lancet 364, 937-952.

Dipl.-Psych. Dr. phil. Budde, Hans-Günter
Drosselweg 1
55583 Bad Münster am Stein-Ebernburg
Telefon 06708 1517
Fax 06708 64185
E-mail heide.dr.budde@t-online.de

Psychologischer Psychotherapeut / Verhaltenstherapie
Spezielle Schmerzpsychotherapie
Medizinische Hypnose

